

QUANTUM $\mathfrak{sl}(n, \mathbb{C})$ LINK INVARIANTS

MYEONG-JU JEONG AND DONGSEOK KIM

ABSTRACT. Kuperberg generalized Temperley-Lieb algebras to the $\mathfrak{sl}(3)$ web spaces, we extend it to $\mathfrak{sl}(n)$ for $n \geq 4$. We find that the HOMFLY polynomial $P_n(q)$ can be computed by a linear expansion with respect to the relations of the quantum $\mathfrak{sl}(n)$ representation theory. We provide criterions which determine the periodicity of links by using the quantum $\mathfrak{sl}(n)$ HOMFLY polynomial.

1. INTRODUCTION

The HOMFLY polynomial $P_n(q)$ can be calculated uniquely by the following skein relations:

$$\begin{aligned} P_n(\emptyset) &= 1, \\ P_n(\bigcirc \cup D) &= \left(\frac{q^{\frac{n}{2}} - q^{-\frac{n}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}} \right) P_n(D), \\ q^{\frac{n}{2}} P_n(L_+) - q^{-\frac{n}{2}} P_n(L_-) &= (q^{\frac{1}{2}} - q^{-\frac{1}{2}}) P_n(L_0), \end{aligned}$$

where $\emptyset$ is the empty diagram, $\bigcirc$ is the trivial knot and L_+ , L_- and L_0 are three diagrams which are identical except at one crossing as in Figure 1 [14]. For each positive integer n , the HOMFLY polynomial of links specializes to a one variable polynomial that can be recovered from the quantum $\mathfrak{sl}(n)$ representation theory. For $n = 0$, we use $\mathfrak{gl}(1|1)$ to find that $P_0(q)$ is the Alexander polynomial [7]. For $n = 1$, one can see that $P_1(q) = 1$ for all links. For $n = 2$, $P_2(q)$ is the Jones polynomial [5, 18, 19]. The polynomial $P_n(q)$ can be computed by linearly expanding each crossing into a sum of planar cubic graphs as in Figure 2 [14]. The edges of these planar cubic graphs are oriented and colored by 1 or 2.



In the language of the representation theory [9, 11], the color 1 of an edge presents the vector representation V of the quantum $\mathfrak{sl}(n)$, and the color 2 for its exterior power $\wedge^2 V$. The trivalent vertex is the unique (up to scaling) intertwiner of $V^{\otimes 2} \otimes \wedge^2 V$. This setup works for arbitrary exterior powers of V : oriented edges of graphs carry colors from 1 to $n - 1$ that denote the fundamental representations of the quantum $\mathfrak{sl}(n)$ [14]. Kuperberg generalized Temperley-Lieb algebras to $\mathfrak{sl}(3)$ web spaces [9]. In section 2 we develop the quantum $\mathfrak{sl}(n)$ representation theory by extending the idea of webs in [14]. Sikora also found a generalization of webs to the quantum $\mathfrak{sl}(n)$ [20]. By expanding all crossings as in Figures 17 and 18 Murakami, Ohtsuki and Yamada found a regular isotopy invariant

Date: June 20, 2005, revised March 1, 2006.

2000 Mathematics Subject Classification. Primary 57M27; Secondary 57M25, 57R56.

Key words and phrases. HOMFLY polynomial, Representation theory, Quantum special linear algebras.

The first author was supported in part by the Post-doctoral Fellowship Program of KOSEF and the second author was supported in part by KRF Grant M02-2004-000-20044-0.

FIGURE 1. The shape of L_+ , L_- and L_0 .FIGURE 2. Expansions of crossings for $P_n(q)$.

$[D]_n$ [14]. In section 3 we modify writhe suitably to define an isotopy invariant $K_n(L, \mu)$ where μ is a coloring of L . Using the quantum $\mathfrak{sl}(n)$ representation theory, we show $K_n(L, \mu)$ can be computed by a linear expansion with respect to the relations of web spaces in Theorem 2.4.

A link L in S^3 is p -periodic if there exists a periodic homeomorphism h of order p such that $\text{fix}(h) \cong S^1$, $h(L) = L$ and $\text{fix}(h) \cap L = \emptyset$ where $\text{fix}(h)$ is the set of all fixed points of h . It is well known that if we consider S^3 as $\mathbb{R} \cup \{\infty\}$, we can assume that h is a rotation by $2\pi/p$ angle around the z -axis. If L is a periodic link, we denote its factor link by $\overline{L}$. Murasugi [12] found a strong relation between the Alexander polynomials of a periodic link and its factor link. Murasugi also found a similar relation for the Jones polynomials of L and $\overline{L}$ [13]. Park and the first author gave a criterion for periodic knots and lens knots by using Vassiliev invariants [6]. There are various result to decide the periodicity of links [15, 16, 21–24]. Chbili generalized it to $P_3(q)$ using the quantum $\mathfrak{sl}(3)$ representation theory and stated a conjecture that the same statement is true for $P_n(q)$ where $n \geq 2$ [1]. In this paper we find a positive answer after a slight modification.

Theorem 1.1. *Let p be a positive integer and L be a p -periodic link in S^3 with its factor link $\overline{L}$. Let μ be a p -periodic coloring of L and $\overline{\mu}$ be the induced coloring of $\overline{L}$. Then for $n \geq 0$,*

$$K_n(L, \mu) \cong K_n(\overline{L}, \overline{\mu})^p \quad \text{modulo } \mathcal{I}_n,$$

where $\mathcal{I}_n$ is the ideal of $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ generated by p and $\begin{bmatrix} n \\ i \end{bmatrix}^p - \begin{bmatrix} n \\ i \end{bmatrix}$ for $i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$.


 FIGURE 3. Generators of the quantum $\mathfrak{sl}(n)$ web space.

The quantum integers are defined as

$$\begin{aligned} [n] &= \frac{q^{\frac{n}{2}} - q^{-\frac{n}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}, \\ [n]! &= [n][n-1] \dots [2][1], \\ \begin{bmatrix} n \\ k \end{bmatrix} &= \frac{[n]!}{[n-k]![k]!}. \end{aligned}$$

Let us remark that even if we replace $K_n(L, \mu)$ by $P_n(q)$ in Theorem 1.1, the ideal $\mathcal{I}_n$ remains the same. Precisely, when we have a disjoint union of unknots after expanding crossings and applying relations, the colors of unknots may not be identically 1 in general. Chen and Le found a similar result using a different method [2], the improvements in Theorem 1.1 are the followings: the result is for an isotopy link invariant, p does not have to be prime and the ideal runs only on a finite set, the set of fundamental representations instead of an infinite set, the set of all finite dimensional irreducible representations of the quantum $\mathfrak{sl}(n)$.

We also give a criterion for periodic links by using the invariants K_n of a link and its mirror image in the following theorem.

Theorem 1.2. *Let L be a p -periodic link and let L^* be the link obtained from the mirror image of a diagram of L . Let μ be a coloring of L and μ^* be the coloring of L^* induced from the coloring μ of L . Then we have*

$$K_n(L, \mu) \equiv K_n(L^*, \mu^*) \pmod{(p, q^p - 1)}.$$

In section 2, we develop the quantum $\mathfrak{sl}(n)$ representation theory. In section 3 we prove main theorems and extend Theorem 1.1 to the colored $\mathfrak{sl}(n)$ HOMFLY polynomial.

2. THE QUANTUM $\mathfrak{sl}(n)$ REPRESENTATION THEORY

We refer to [4, 9–11] for the general representation theory. The webs are generated by the two types of shapes in Figure 3 where $i + j + k = n$. At the second web in Figure 3, by using the duality, $(V_{\lambda_i})^* \cong V_{\lambda_{n-i}}$, we can change the directions of the edges to inward and the colors of the resulting edges to $n - i, n - j$ and $n - k$. After this process, we say a vertex has a *sign type* $+$ if the sum of the colors is n , $-$ if the sum is $2n$. The generators in Figure 3 have sign type $+, -$ from left. Naturally we can assign a sign type for each face. Using relations 2.3 and 2.4 in Figure 4, we only need to look at the face of a *valid sign type* in which $+, -$ are appearing alternatively. One can easily see that none of faces with odd numbers of edges have a valid sign type.

A few computations of tensor products of fundamental representations show the relations in Figure 4 (up to scaling): the quantum dimension of the fundamental representation V_{λ_i} is

$$(2.1) \quad \bigcirc^i = \begin{bmatrix} n \\ i \end{bmatrix}$$

$$(2.2) \quad i \xrightarrow{i-j} \begin{array}{c} \xrightarrow{j} \\ \xleftarrow{i-j} \end{array} i = \begin{bmatrix} i \\ j \end{bmatrix} i \longrightarrow i$$

$$(2.3) \quad \begin{array}{c} i \searrow \quad \nearrow i+j+k \\ \quad \nearrow j+k \\ j \nearrow \quad \searrow k \end{array} = \begin{array}{c} i \searrow \quad \nearrow i+j+k \\ \quad \nearrow i+j \\ j \nearrow \quad \searrow k \end{array}$$

$$(2.4) \quad \begin{array}{c} i \searrow \quad \nearrow i+j+k \\ \quad \nearrow j+k \\ j \nearrow \quad \searrow k \end{array} = \begin{array}{c} i \searrow \quad \nearrow i+j+k \\ \quad \nearrow i+j \\ j \nearrow \quad \searrow k \end{array}$$

FIGURE 4. Relations obtained by the quantum dimension and scaling.

$$\{ \begin{array}{c} i \searrow \quad \nearrow j \\ \quad \nearrow l \\ l+j \nearrow \quad \searrow i-l \\ \quad \nearrow l \\ i \nearrow \quad \searrow j \end{array} \}_{l=0}^i \quad \text{or} \quad \{ \begin{array}{c} i \searrow \quad \nearrow j \\ \quad \nearrow l \\ i+l \nearrow \quad \searrow j-l \\ \quad \nearrow l \\ i \nearrow \quad \searrow j \end{array} \}_{l=0}^i$$

FIGURE 5. Two basis webs with the boundary $V_{\lambda_i} \otimes V_{\lambda_j} \otimes V_{\lambda_i}^* \otimes V_{\lambda_j}^*$.

$\begin{bmatrix} n \\ i \end{bmatrix}$, the dimension of the invariant space of $V_{\lambda_i} \otimes V_{\lambda_i}^*$ is 1 and the dimension of the invariant space of $V_{\lambda_i} \otimes V_{\lambda_j} \otimes V_{\lambda_k} \otimes V_{\lambda_i+\lambda_j+\lambda_k}^*$ is 1.

2.1. Rectangular Relations. To discuss rectangular relations, we first prove the following lemma.

Lemma 2.1. (1) *If $i \leq j \leq n - j - 1$, then*

$$\dim(\text{inv}(V_{\lambda_i} \otimes V_{\lambda_j} \otimes V_{\lambda_i}^* \otimes V_{\lambda_j}^*)) = i + 1.$$

(2) *If $j \geq i \geq k \geq 1, n - j - 1 \geq i$ and $n - i - j - 1 \geq l \geq 1$, then*

$$\dim(\text{inv}(V_{\lambda_i} \otimes V_{\lambda_{j+l}} \otimes V_{\lambda_{i+l}}^* \otimes V_{\lambda_j}^*)) = i + 1.$$

Proof. If $i \leq j \leq n - j - 1$, we obtain the following isomorphism by the Clebsch-Gordan formula.

$$V_{\lambda_i} \otimes V_{\lambda_j} \cong V_{\lambda_i+\lambda_j} \oplus V_{\lambda_{i-1}+\lambda_{j+1}} \oplus \cdots \oplus V_{\lambda_1+\lambda_{i+j}} \oplus V_{\lambda_{i+j+1}}.$$

$$\begin{aligned}
 (2.5) \quad & \begin{array}{c} i \quad j+k-i \quad j+l \\ \swarrow \quad \rightarrow \quad \nearrow \\ j+k \quad i-k+l \quad j+l \\ \nwarrow \quad \leftarrow \quad \searrow \\ j \quad k \quad i+l \end{array} = \sum_{m=0}^i \begin{bmatrix} l \\ k-m \end{bmatrix} \begin{array}{c} i \quad m \quad j+l \\ \swarrow \quad \rightarrow \quad \nearrow \\ i-m \quad j+l+m \quad j+l \\ \nwarrow \quad \leftarrow \quad \searrow \\ j \quad j-i+m \quad i+l \end{array} \\
 (2.6) \quad & \begin{array}{c} i \quad k \quad j+l \\ \swarrow \quad \rightarrow \quad \nearrow \\ i+k \quad j-k+l \quad j+l \\ \nwarrow \quad \leftarrow \quad \searrow \\ j \quad i-j+k \quad i+l \end{array} = \sum_{m=0}^j \begin{bmatrix} l \\ k-m \end{bmatrix} \begin{array}{c} i \quad i-j+m \quad j+l \\ \swarrow \quad \rightarrow \quad \nearrow \\ j-m \quad i+l+m \quad j+l \\ \nwarrow \quad \leftarrow \quad \searrow \\ j \quad m \quad i+l \end{array}
 \end{aligned}$$

FIGURE 6. Equations in Lemma 2.2.

$$\begin{array}{c} i \quad j+l \\ \swarrow \quad \nearrow \\ j-i \quad j+l \\ \nwarrow \quad \nearrow \\ j \quad i+l \end{array} = \begin{bmatrix} l \\ 0 \end{bmatrix} \begin{array}{c} i \quad j+l \\ \swarrow \quad \nearrow \\ j-i \quad j+l \\ \nwarrow \quad \nearrow \\ j \quad i+l \end{array}$$

 FIGURE 7. Initial induction step $k = 0$ and $j \geq i$.

$$\begin{array}{c} j+l \\ \swarrow \quad \nearrow \\ j+k \quad l-k \\ \nwarrow \quad \nearrow \\ j \quad k \quad l \end{array} = \begin{array}{c} j+l \\ \swarrow \quad \nearrow \\ j \quad l-k \\ \nwarrow \quad \nearrow \\ j \quad k \quad l \end{array} = \begin{bmatrix} l \\ k \end{bmatrix} \begin{array}{c} j+l \\ \swarrow \quad \nearrow \\ j \quad l \\ \nwarrow \quad \nearrow \\ j \quad l \end{array}$$

 FIGURE 8. Initial induction step $i = 0$.

For irreducible representations V, W of a simple Lie algebra, by a simple application of Schur's lemma we find

$$\dim(\text{Inv}(V \otimes W^*)) = \begin{cases} 1 & \text{if } V \cong W, \\ 0 & \text{if } V \not\cong W. \end{cases}$$

These two facts imply that $\dim(\text{inv}(V_{\lambda_i} \otimes V_{\lambda_j} \otimes V_{\lambda_i}^* \otimes V_{\lambda_j}^*)) = i + 1$. Similarly one can prove the other. $\square$

From Lemma 2.1, we know the number of basis webs that we need for each expansion. For $n - i - 1 \geq j \geq i \geq 0$, we can have two sets of basis webs and each has the same sign type as in Figure 5. Throughout the section we will use the basis in the left hand side of Figure 5. There are only two possible types of rectangular relations as in Lemma 2.2, all other can be taken care of by relations 2.3 and 2.4 in Figure 4. The equation 2.5 in Lemma 2.2 was appeared in [14] without a proof.

Lemma 2.2. *For $n - i - 1 \geq j \geq i \geq k \geq 0$ and $n - i - j - 1 \geq l \geq 0$, we find equation 2.5 in Figure 6. For $n - j - 1 \geq i \geq j \geq k \geq 1$ and $n - i - j - 1 \geq l \geq 1$, we find equation 2.6 in*




FIGURE 9.

$$\begin{aligned}
 & \begin{array}{c} i-1 \quad j+l+1 \\ \swarrow \quad \searrow \\ \text{web} \\ \swarrow \quad \searrow \\ j \quad i+l \end{array} = \begin{bmatrix} j+k-i+1 \\ 1 \end{bmatrix} \begin{array}{c} i-1 \quad j+l+1 \\ \swarrow \quad \searrow \\ \text{web} \\ \swarrow \quad \searrow \\ j \quad i+l \end{array} \\
 &= \begin{bmatrix} j+k-i+1 \\ 1 \end{bmatrix} \sum_{m=0}^{i-1} \begin{bmatrix} l+1 \\ k-m \end{bmatrix} \begin{array}{c} i-1 \quad j+l+1 \\ \swarrow \quad \searrow \\ \text{web} \\ \swarrow \quad \searrow \\ j \quad i+l \end{array}
 \end{aligned}$$

FIGURE 10.

Figure 6. Just for these two equations, we use a different convention of quantum integers that $\begin{bmatrix} 0 \\ 0 \end{bmatrix} = 1$ but $\begin{bmatrix} 0 \\ k \end{bmatrix} = 0$ if $k \neq 0$.

Proof. Let $a(i, m)$ be the coefficients in the righthand side of the equations. We induct on $(\min\{i, j\}, k)$ in lexicographic order. The key idea is to prove both equations simultaneously. If $k = 0$ and $j \geq i$, we find the equation in Figure 7. For the case $k = 0$ and $i \geq j$, it is identical except the weight on the horizontal arrow is replaced by the weight $i - j$ of the opposite direction. For $i = 0$, we find the equations in Figure 8. One can do for the case $j = 0$ similarly. $\square$

Now we proceed to the induction step. Let us look at the case $n - i - 1 \geq j \geq i \geq k \geq 0$, $n - i - j - 1 \geq l \geq 0$. On the top of each web in equations 2.5 and 2.6 in Figure 6, we can attach i different H 's, given in Figure 9 where $i \geq s \geq 1$. If $s = i$, we can easily get

$$\begin{bmatrix} j+k \\ i \end{bmatrix} \begin{bmatrix} i+l \\ k \end{bmatrix} = \sum_{m=0}^i a(i, m) \begin{bmatrix} j+l+m \\ m \end{bmatrix} \begin{bmatrix} j \\ i-m \end{bmatrix}.$$

From the case $s = 1$, we find that the left hand side of equation 2.6 can be changed as the equations in Figure 10. We use the induction hypothesis to find that each term in right hand side of equation 2.6 can be transformed as the equations in Figure 11 where $\alpha = [-i + j + l + m + 1]$, $\beta = 1$ and $\gamma = [-i + j + m + 1]$. By comparing coefficients of each basis element, we get i equations




FIGURE 11.

$$[-i + j + k + 1] \begin{bmatrix} l + 1 \\ k - t \end{bmatrix} = a(i, t)[-i + j + t + 1] + a(i, t + 1)[-i + j + l + t + 2],$$

where $i - 1 \geq t \geq 0$. Since these $i + 1$ equations are independent, we plug in the answer to equations to check these are correct. One can follow the proof for the second case.

2.2. The quantum $\mathfrak{sl}(n)$ skein modules. The authors have been trying to find a complete relation of the quantum of $\mathfrak{sl}(n)$ representation theory, but we find that the bigger n , the bigger the size of a polygon we have to find a suitable relation: there are a rectangular relation for the quantum of $\mathfrak{sl}(3)$, a hexagonal relation for the quantum of $\mathfrak{sl}(4)$ and an octagonal relation for the quantum of $\mathfrak{sl}(6)$. In particular, it was conjectured a complete set of relations for the quantum $\mathfrak{sl}(4)$ representation theory [8]. For large n , we find the following remark.

Remark 2.3. *We conjecture that any $2n$ polygon of the sign type $(+, -, +, \dots, -)$ can be expanded to a sum of webs of polygons of smaller sizes and $2n$ polygons of the sign type $(-, +, -, \dots, +)$.*

Somehow the study of the quantum $\mathfrak{sl}(n)$ representation theory is not complete yet, but we can prove the following theorem.

Theorem 2.4. *The quantum $\mathfrak{sl}(n)$ skein module of the plane or sphere has dimension 1.*

Proof. If we look at these webs without decorations, they are directed, weighted, trivalent planar graphs. We will consider these graphs are on $\mathbb{S}^2$ instead of $\mathbb{R}^2$. Let V, E and F be the number of vertices, edges and faces in a graph on $\mathbb{S}^2$, respectively. Let F_i be the number of faces with i edges. We can easily find the following equations:

$$\begin{aligned}
(2.7) \quad & 3V = 2E \\
& 2 = V - E + F \\
& 2E = \sum_i iF_i
\end{aligned}$$

Throughout the section all webs are assumed to be without boundary unless we state differently. We will show the dimension of the space of webs without boundary is 1. Suppose it is not true, then there exists a web which is not an element in $\mathbb{C}[q^{\frac{1}{2}}, q^{-\frac{1}{2}}]$, say D . We assume that D has the smallest number of faces among all counterexamples. Since a face of size less than 4 can be removed by using relations we found in Figure 4, we will assume that all faces in D have at least 4 edges and we further assume that D has the maximum number of rectangular faces.

Lemma 2.5. *If there exists a pentagonal face in D , it is adjacent to a unique rectangular face.*

Proof. It is easy to see that there are no two adjacent rectangular faces by considering Figure 5. Thus if a pentagon is not adjacent to a rectangle, the sizes of all adjacent faces of the pentagon are bigger than 5 because an adjoint pentagon can be changed to a rectangle. By applying relations 2.3 and 2.4 in Figure 4 to the pentagonal face, it can be changed to a rectangle without changing the total number of faces in D but increasing the number of the rectangles by 1. This contradicts the maximality of the number of the rectangular faces. Therefore, it must be adjacent to a rectangle. There are five possible shapes which can be obtained from the web in Figure 12 by $\frac{2\pi}{5}$ rotation. By applying equations in Lemma 2.2, one can see that any two out of these five shapes are related as the equation in Figure 13. Therefore, the adjacent rectangle is unique. $\square$

A rectangular face is called *isolated* if the sizes of all adjacent faces (after making valid sign types) are bigger than 5. Otherwise, it is adjacent to the unique pentagon by Lemma 2.5. Now we will look at the neighborhood of each isolated rectangular face in subsection 2.2.1 and obtain the following inequality.

$$\begin{aligned}
4F_4 &\leq 3F_7 + 4F_8 + 4F_9 + 5F_{10} + 5F_{11} + \dots + \lfloor \frac{n}{2} \rfloor F_n + \dots, \\
5F_4 &\leq 4F_7 + 5F_8 + 5F_9 + 7F_{10} + 7F_{11} + \dots + nF_n + \dots,
\end{aligned}$$

for $n \geq 12$.

By looking at the neighborhood of a pentagon adjacent to a unique rectangle, we find the following inequality in subsection 2.2.2.

$$5(F_5 + F_4) \leq 4F_8 + 4F_9 + 5F_{10} + 5F_{11} + \dots + \lfloor \frac{n}{2} \rfloor F_n + \dots$$

Now let us proceed to the proof. By combining these two inequalities and the inequality, $(n-6) \geq \frac{2}{5}n$ for all $n \geq 12$, we get the following inequality.




FIGURE 12. A valid shapes of a pentagon face.




FIGURE 13. Pentagonal relations.

$$\begin{aligned}
 5F_5 + 10F_4 &\leq 3F_7 + 9F_8 + 9F_9 + 12F_{10} + 12F_{11} \\
 &\quad + \dots + (n + \lfloor \frac{n}{2} \rfloor)F_n + \dots, \\
 2F_4 + F_5 &\leq 1F_7 + 2F_8 + 3F_9 + 4F_{10} + 5F_{11} \\
 &\quad + \dots + (n - 6)F_n + \dots
 \end{aligned}
 \tag{2.8}$$

Then,

$$\begin{aligned}
 6F &= 6F_4 + 6F_5 + 6F_6 + 6F_7 + 6F_8 + \dots + 6F_n + \dots \\
 &\leq 4F_4 + 5F_5 + 6F_6 + 7F_7 + 8F_8 + \dots + nF_n + \dots = 2E.
 \end{aligned}$$

If we plug this inequality into equations 2.7, we find a contradiction as

$$2 = V - E + F \leq \frac{2}{3}E - E + \frac{1}{3}E = 0.$$

Therefore, this completes the proof. □

2.2.1. Isolated rectangular faces. From Lemma 2.5, we know all adjacent faces are either hexagons or bigger polygons. Moreover, all adjacent hexagonal faces must have a valid sign type (otherwise, they become two adjacent rectangular faces). We will divide cases with the number of adjacent hexagonal faces. If there is no adjacent hexagonal face, then it is surrounded by at least four faces of sizes bigger than or equal to 8 and

$$4F_4 \leq 4F_8 + 4F_9 + \dots + \lfloor \frac{n}{2} \rfloor F_n + \dots$$




FIGURE 14. Just one hexagon in the neighborhood of the rectangle.




FIGURE 15. Two non-adjacent hexagons in the neighborhood of the rectangle.

If there is just one hexagonal face, Lemma 2.2 allows us to change a rectangular face to the sum of rectangular (or less) faces with the opposite sign type, this forces a change which is called a *swapping* of the rectangle and the hexagon as in Figure 14 without any restrictions on the sizes of neighbor faces. If there is only one hexagon in the neighborhood of the rectangle after a swapping, then we find the desired inequality as the sizes of faces indicated by numbers in Figure 14. If there are more than one hexagon, it will be dealt in the next cases.

If there are two adjacent hexagonal faces, there are also two possibilities: If these two hexagons are adjacent, the rectangular face can be changed into two rectangles which contradicts the hypothesis that D has the maximum number of rectangles as in Figure 16. If these two hexagons are not adjacent each other, then we can continue to swap and we repeat the process at the new rectangle. Since there are only finitely many faces, it stops at finite steps. By tracing the swap moves, we find that it is just like in Figure 15. For all other cases, we must have at least two adjacent hexagons. However, it can be expanded into a sum of webs which has either less faces or more rectangular faces as shown in Figure 16 which contradict the hypothesis of D .

As we have seen, for each rectangular face, there are at least four polygons which have more than seven faces. It is easy to see that for each face of size $n \geq 7$, there are at most $\frac{n}{2}$ (or $\frac{n-3}{2}$) rectangular faces which can travel by swapping moves to the given face if n is even (or odd, respectively). Therefore, we get the inequality we desired.

2.2.2. Pentagonal face with a unique adjacent rectangular faces. We will treat these two adjacent polygons as a pentagon in this subsection.

Lemma 2.6. *The size of all adjacent polygons must be bigger than 7.*




FIGURE 16. Two adjacent hexagons in the neighborhood of the rectangle.

Proof. First we already know these polygons can be neither rectangles nor pentagons. Suppose one of them is either a hexagon or a heptagon. Either case, we transform the shapes of these polygons that the rectangular face is adjacent to the hexagon (now rectangle has a valid sign type, we can change the heptagon to a hexagon). By the relation in Figure 13, we assume the hexagonal face is adjacent to the rectangular face. Then as we have seen in Figure 14, the rectangle can be separated from the pentagon. Then the remaining pentagon can be changed to a rectangle which contradict the maximality of the number of rectangles in D . $\square$

One can easily find the following inequality.

$$5(F_5 + F_4) \leq 4F_8 + 4F_9 + 5F_{10} + 5F_{11} + \dots + \lfloor \frac{n}{2} \rfloor F_n + \dots$$

3. LINK INVARIANTS

When links are decorated by the fundamental representations V_{λ_i} of the quantum $\mathfrak{sl}(n)$, denoted by i , Murakami, Ohtsuki and Yamada [14] found a quantum invariant for framed links by resolving each crossing in a link diagram D of L as shown in Figures 17 and 18.

For negative crossings, we replace q with q^{-1} . $P_n(q)$ is the special case of $q^{-\omega(D)\frac{n}{2}}[D]_n$, when all components are colored by the fundamental representation V_{λ_1} . They showed that $P_n(q)$ is an isotopy invariant and that $[D]_n$ is a regular isotopy invariant for other colorings [14]. However, one can make it a link invariant by using a suitable writhe but we have to be careful since there are more than one colors. For a coloring μ of a diagram D of a link L , we first consider a *colored writhe* $\omega_i(D)$ as the sum of writhes of components colored by i . Then we set

$$K_n(L, \mu) = \prod_i q^{-\omega_i(D)\frac{i(n-i+1)}{2}} [D]_n,$$

where the product runs over all colors i .

Theorem 3.1. $K_n(L, \mu)$ is an isotopy invariant of a link L , i.e., invariant under Reidemeister moves I, II and III.

Proof. By Theorem 2.4, we know $K_n(L, \mu)$ is well defined. It was shown that $[D]_n$ is a regular isotopy invariant [14], so is $K_n(L, \mu)$. One can find the equations in Figure 19 by

$$\left[\begin{array}{c} i \swarrow \nearrow j \\ j \nearrow \swarrow i \end{array} \right]_n = \sum_{k=0}^i (-1)^{k+(j+1)i} q^{\frac{(i-k)}{2}} \left[\begin{array}{c} i \swarrow \nearrow j \\ j+k \nearrow \swarrow i-k \\ j \nearrow \swarrow i \end{array} \right]_n$$

FIGURE 17. Skein expansion case 1 : $j \geq i$.

$$\left[\begin{array}{c} i \swarrow \nearrow j \\ j \nearrow \swarrow i \end{array} \right]_n = \sum_{k=0}^j (-1)^{k+(i+1)j} q^{\frac{(j-k)}{2}} \left[\begin{array}{c} i \swarrow \nearrow j \\ j-k \nearrow \swarrow i+k \\ j \nearrow \swarrow i \end{array} \right]_n$$

FIGURE 18. Skein expansion case 2 : $j < i$.

$$\left[\begin{array}{c} i \swarrow \nearrow \\ \nearrow \swarrow \end{array} \right]_n = q^{\frac{i(n-i+1)}{2}} \left[\begin{array}{c} i \swarrow \\ \nearrow \end{array} \right]_n, \quad \left[\begin{array}{c} i \swarrow \nearrow \\ \nearrow \swarrow \end{array} \right]_n = q^{-\frac{i(n-i+1)}{2}} \left[\begin{array}{c} i \swarrow \\ \nearrow \end{array} \right]_n$$

FIGURE 19. Resolving Reidemeister move I.

relation 2.2 and a routine induction. Then $K_n(L, \mu)$ is invariant under Reidemeister move I. $\square$

3.1. Proof of Theorem 1.1. Let μ be a p -periodic coloring of L . Let R be the fundamental region by the action h . Let $\mathcal{C}$ be the set of all crossings of L and let $\overline{\mathcal{C}}$ be the set of all crossings in the region R . Let $i(c)$ and $j(c)$ be the weights of two components at the crossing c as in Figures 17 and 18. Let $J(c)$ be the minimum of $i(c)$ and $j(c)$ for the crossing c . Let D be the diagram after the expansion by the equations in Figures 17 and 18. Let D' be the diagram obtained by identical expanding of the crossings which are the same by the action h and D'' be the diagram obtained from D' by identical applications of relations at the faces which are in the same orbit by the action h . If any expansion of crossings occurs in R , it must be used identically for all other $p-1$ copies of R . Otherwise there will be p identical shapes by the rotation of order p , then it is congruent to zero modulo p . This implies the first congruence in equation 3.1. By the same philosophy, if any expansion of relations occurs in R , it must be used identically for all other $p-1$ copies of R . Otherwise it is congruent to zero modulo p and this implies the second congruence. Let us remark that we have not used relations that might occur in the faces which are not entirely contained in R . If there is an unknot in $\overline{D''}$ which was in D'' , there are p copies in D'' . Thus raising p -power gives us the equality. But if there is an unknot in $\overline{D''}$ which was not in D'' , it should be counted once in D'' and $\overline{D''}$, thus we have to use $\mathcal{I}_n$ congruence.

$$\begin{aligned}
(3.1) \quad K_n(L, \mu) &= \prod_{c \in \mathcal{C}} \sum_{k=0}^{J(c)} (-1)^{k+(i(c)+1)j(c)} q^{\frac{(j(c)-k)}{2}} [D]_n \\
&\equiv \prod_{\bar{c} \in \bar{\mathcal{C}}} \sum_{k=0}^{J(\bar{c})} ((-1)^{k+(i(\bar{c})+1)j(\bar{c})} q^{\frac{(j(\bar{c})-k)}{2}})^p [D']_n \quad (\text{mod } p) \\
&\equiv \prod_{\bar{c} \in \bar{\mathcal{C}}} \sum_{k=0}^{J(\bar{c})} ((-1)^{k+(i(\bar{c})+1)j(\bar{c})} q^{\frac{(j(\bar{c})-k)}{2}})^p [D'']_n \quad (\text{mod } p) \\
&\equiv \prod_{\bar{c} \in \bar{\mathcal{C}}} \sum_{k=0}^{J(\bar{c})} ((-1)^{k+(i(\bar{c})+1)j(\bar{c})} q^{\frac{(j(\bar{c})-k)}{2}})^p ([\overline{D''}]_n)^p \quad (\text{mod } \mathcal{I}_n)
\end{aligned}$$

where $\overline{D''} = D''/\mathbb{Z}_p$, $\mathbb{Z}_p$ is generated by the action h , and $\mathcal{I}_n$ is the ideal of $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ generated by p and $\begin{bmatrix} n \\ i \end{bmatrix}^p - \begin{bmatrix} n \\ i \end{bmatrix}$ for $i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$. The following congruences should be straightforward.

$$\begin{aligned}
&\equiv \prod_{\bar{c} \in \bar{\mathcal{C}}} \sum_{k=0}^{J(\bar{c})} ((-1)^{k+(i(\bar{c})+1)j(\bar{c})} q^{\frac{(j(\bar{c})-k)}{2}} [\overline{D}]_n)^p \quad (\text{mod } p) \\
&\equiv \left(\prod_{\bar{c} \in \bar{\mathcal{C}}} \sum_{k=0}^{J(\bar{c})} (-1)^{k+(i(\bar{c})+1)j(\bar{c})} q^{\frac{(j(\bar{c})-k)}{2}} [\overline{D}]_n \right)^p \quad (\text{mod } p) \\
&\equiv (K_n(\overline{L}, \overline{\mu}))^p \quad (\text{mod } p)
\end{aligned}$$

Next, we will extend Theorem 1.1 to the colored $\mathfrak{sl}(n)$ HOMFLY polynomial of a periodic link whose components are colored by irreducible representations of quantum $\mathfrak{sl}(n)$. For a given colored link L of l components say, $L_1, L_2, \dots, L_l$, where each component L_i is colored by an irreducible representation $V_{a(i)_1 \lambda_1 + a(i)_2 \lambda_2 + \dots + a(i)_{n-1} \lambda_{n-1}}$ of $\mathfrak{sl}(n)$ and $\lambda_1, \lambda_2, \dots, \lambda_{n-1}$ are the fundamental weights of $\mathfrak{sl}(n)$. The coloring is denoted by $\mu = (a(1)_1 \lambda_1 + a(1)_2 \lambda_2 + \dots + a(1)_{n-1} \lambda_{n-1}, a(2)_1 \lambda_1 + a(2)_2 \lambda_2 + \dots + a(2)_{n-1} \lambda_{n-1}, \dots, a(l)_1 \lambda_1 + a(l)_2 \lambda_2 + \dots + a(l)_{n-1} \lambda_{n-1})$. First we replace each component L_i by $a(i)_1 + a(i)_2 + \dots + a(i)_{n-1}$ copies of parallel lines and each $a(i)_j$ line is colored by the weight λ_j . Then we put a clasp of weight $(a(i)_1 \lambda_1 + a(i)_2 \lambda_2 + \dots + a(i)_{n-1} \lambda_{n-1})$ for L_i . If we assume the clasps are far away from crossings, we expand each crossing as in Figure 17 and 18, then clasps [25]. The value we find after removing all faces by using the relations is the colored $\mathfrak{sl}(n)$ HOMFLY polynomial $G_n(L, \mu)$ of L .

Theorem 3.2. *Let p be a positive integer and L be a p -periodic link in S^3 with the factor link $\overline{L}$. Let μ be a p -periodic coloring of L and $\overline{\mu}$ be the induced coloring of $\overline{L}$. Then for $n \geq 2$,*

$$G_n(L, \mu) \cong G_n(\overline{L}, \overline{\mu})^p \quad \text{modulo } \mathcal{I}_n,$$

where $\overline{L}$ is the factor link and $\mathcal{I}_n$ is the ideal of $\mathbb{Z}[q^{\pm\frac{1}{2}}]$ generated by p and $\begin{bmatrix} n \\ i \end{bmatrix}^p - \begin{bmatrix} n \\ i \end{bmatrix}$ for $i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$.

Proof. Since all clasps are idempotents, we put p -copies of clasps to each copy of the fundamental region by the action h of the periodicity of L . Thus without expanding the clasps, we obtain the theorem by the same idea of the proof of the Theorem 1.1. $\square$

3.2. Proof of Theorem 1.2. For a colored link diagram D we denote its mirror image by D^* which is the colored link diagram obtained from D by changing all of the crossings. We study another necessary condition for a colored link to be periodic by using the invariant $K_n(L, q)$. Let L be a colored link diagram with a crossing x . Let L_+ , L_- , and L_0 be the link diagrams obtained by resolving the crossing x as shown in Figure 4 respectively. Then from the Figure 17, 18, we see that there exist web diagrams $L_1, \dots, L_m$ and polynomials $f_1, \dots, f_m \in \mathbb{Z}[q^{\pm\frac{1}{2}}]$ for some positive integer m such that

$$[L_+]_n - [L_-]_n = (q^{\frac{1}{2}} - q^{-\frac{1}{2}})(f_1[L_1]_n + \dots + f_m[L_m]_n).$$

In particular, by applying this relation repeatedly for a colored periodic link we obtain the following lemma.

Lemma 3.3. *For a colored p -periodic link diagram D and its mirror image D^* , we have*

$$[D]_n \equiv [D^*]_n \pmod{(p, q^p - 1)}.$$

Proof. Let π be the quotient map of h and $\overline{D}$ be the factor link of D . Then $\pi^{-1}(\overline{D}) = D$. We fix a crossing x for a nontrivial diagram $\overline{D}$, and let $\overline{D}_+$ and $\overline{D}_-$ be the colored link diagram obtained by changing the crossing x to a positive crossing and a negative crossing respectively.

Let $\overline{D}_1, \dots, \overline{D}_m$ be web diagrams and let $f_1, \dots, f_m \in \mathbb{Z}[q^{\pm\frac{1}{2}}]$ be polynomials for some positive integer m such that

$$[\overline{D}_+]_n - [\overline{D}_-]_n = (q^{\frac{1}{2}} - q^{-\frac{1}{2}})(f_1[\overline{D}_1]_n + \dots + f_m[\overline{D}_m]_n).$$

Then by considering the periodic action induced from the map h , we find the following congruence relation.

$$\begin{aligned} & [\pi^{-1}(\overline{D}_+)]_n - [\pi^{-1}(\overline{D}_-)]_n \\ & \equiv (q^{\frac{1}{2}} - q^{-\frac{1}{2}})^p (f_1^p[\pi^{-1}(\overline{D}_1)]_n + \dots + f_m^p[\pi^{-1}(\overline{D}_m)]_n) \pmod{(p)} \\ & \equiv 0 \pmod{(p, q^p - 1)}. \end{aligned}$$

Since $[\pi^{-1}(\overline{D}_+)]_n \equiv [\pi^{-1}(\overline{D}_-)]_n \pmod{(p, q^p - 1)}$ for each crossing of $\overline{D}$ and since there are finitely many crossings in $\pi^{-1}(\overline{D})$, we see that $[\pi^{-1}(\overline{D})]_n \equiv [\pi^{-1}(\overline{D}^*)]_n \pmod{(p, q^p - 1)}$, where $\overline{D}^*$ is the mirror image of $\overline{D}$. Therefore we have

$$[D]_n \equiv [D^*]_n \pmod{(p, q^p - 1)}.$$

$\square$

Now, we are ready to prove Theorem 3.2. For a tangle T , we denote its closure by $CL(T)$ if it is well defined. Let L be a p -periodic link and T be a tangle such that L is the closure $CL(T^p)$ of the tangle T^p which is p times self-product of T . Let D be a diagram of L and x be a crossing of the diagram D . Let T_+ and T_- be the diagram obtained from the diagram of T by changing the crossing x to a positive crossing and negative crossing respectively. If the two colorings of the strands near the crossing x are different then by using Lemma 3.3, we see that

$$\begin{aligned} & K_n(CL((T_+)^p), \mu) - K_n(CL((T_-)^p), \mu) \\ &= \Pi_i q^{-w_i(D) \frac{i(n-i+1)}{2}} ([CL((T_+)^p)]_n - [CL((T_-)^p)]_n) \\ &\equiv 0 \pmod{(p, q^p - 1)}. \end{aligned}$$

If two colorings of the strands near the crossing x are equal, say i , then $w_i(CL((T_+)^p)) = w_i(CL((T_-)^p)) + 2p$. Then we see that there exist a polynomial $g \in \mathbb{Z}[q^{\pm \frac{1}{2}}]$ such that

$$\begin{aligned} & K_n(CL((T_+)^p), \mu) - K_n(CL((T_-)^p), \mu) \\ &= g([CL((T_+)^p)]_n - q^{pi(n-i+1)} [CL((T_-)^p)]_n) \\ &\equiv 0 \pmod{(p, q^p - 1)}. \end{aligned}$$

The last congruence relation in the above formulae can be obtained by using Lemma 3.3. Thus we see that

$$K_n(CL((T_+)^p), \mu) \equiv K_n(CL((T_-)^p), \mu) \pmod{(p, q^p - 1)}.$$

Since T can be obtained from its mirror image T^* by a finite sequence of crossing changes, we get

$$K_n(L, \mu) \equiv K_n(L^*, \mu^*) \pmod{(p, q^p - 1)}.$$

It completes the proof the theorem.

Acknowledgements

The authors would like to thank Greg Kuperberg for introducing the subject and helpful discussion, Mikhail Khovanov, Jaeun Lee, Hitoshi Murakami, Myungsoo Seo for their attention to this work. Also, the $\text{\TeX}$ macro package PSTricks [17] was essential for typesetting the equations and figures.

REFERENCES

1. N. Chbili, *The quantum $SU(3)$ invariant of links and Murasugi's congruence*, Topology appl., 122, (2002), 479–485.
2. Q. Chen and T. Le, *Quantum invariants and periodic links and periodic manifolds*, preprint, arXiv:math.QA/0408358.
3. I. Frenkel and M. Khovanov, *Canonical bases in tensor products and graphical calculus for $U_q(\mathfrak{sl}_2)$* , Duke Math. J., 87(3), (1997) 409–480.
4. W. Fulton and J. Harris, *Representation theory*, Graduate Texts in Mathematics, 129, Springer-Verlag, New York-Heidelberg-Berlin, 1991.

5. V. F. R. Jones, *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math., 126, (1987), 335–388.
6. M.-J. Jeong and C.-Y. Park, *Lens knots, periodic knots and Vassiliev invariants*, J. of Knot Theory and Its Ramifications, Vol. 13 (2004), 1041–1056.
7. M. Khovanov, private communication.
8. D. Kim, *Graphical Calculus on Representations of Quantum Lie Algebras*, Thesis, UC Davis, 2003, arXiv:math.QA/0310143.
9. G. Kuperberg, *Spiders for rank 2 Lie algebras*, Comm. Math. Phys., 180(1), (1996) 109–151, arXiv:q-alg/9712003.
10. C. Kassel, M. Rosso and V. Turaev, *Quantum groups and knot invariants*, Panoramas et Syntheses, 5, Societe Mathematique de France, 1997.
11. M. Khovanov and L. Rozansky, *Matrix factorizations and link homology*, preprint, arXiv:QA/0401268.
12. K. Murasugi, *On periodic knots*, Comment. Math. Helv., 46 (1971), 162–174.
13. K. Murasugi, *The Jones polynomials of periodic links*, Pacific J. Math., 131 (1988), 319–329.
14. H. Murakami and T. Ohtsuki and S. Yamada, *HOMFLY polynomial via an invariant of colored plane graphs*, L'Enseignement Mathematique, t., 44 (1998), 325–360.
15. J. H. Przytycki, *On Murasugi's and Traczyk's criteria for periodic links*, Math. Ann. 283 (1989), 465–478.
16. J. H. Przytycki and A. Sikora, *SU_n -quantum invariants for periodic links*, Preprint.
17. T. Van Zandt. PSTricks: PostScript macros for generic $\text{\TeX}$. Available at `ftp://ftp.princeton.edu/pub/tvz/`.
18. N. Yu. Reshetikhin and V. G. Turaev, *Ribbob graphs and their invariants derived from quantum groups*, Comm. Math. Phys., 127 (1990), 1–26.
19. N. Yu. Reshetikhin and V. G. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math., 103 (1991), 547–597.
20. A. Sikora., *Skein theory for $SU(n)$ -quantum invariants*, Algebr. Geom. Topol. 5 (2005) 865–897, arXiv:math.QA/0407299.
21. P. Traczyk, *A criterion for knots of period 3*, Topology and its Appl. 36 (1990), 275–281.
22. Y. Yokota, *The skein polynomial of periodic knots*, Math. Ann. 291(2) (1991), 281–291.
23. Y. Yokota, *The Kauffman polynomial of periodic knots*, Topology 32(2) (1993), 309–324.
24. Y. Yokota, *Polynomial invariants of periodic knots*, J. Knot Theory and Its Ramifications 5(4) (1996), 553–567.
25. Y. Yokota, *Skein and quantum $SU(N)$ invariants of 3-manifolds*, Math. Ann., 307 (1997), 109–138.

DEPARTMENT OF MATHEMATICS, KOREAN SCIENCE ACADEMY, PUSAN, 614-822, KOREA
E-mail address: `determiner@hanmail.net`

DEPARTMENT OF MATHEMATICS, COLLEGE OF NATURAL SCIENCE, KYUNGPOOK NATIONAL UNIVERSITY, TAEGU, 702-201, KOREA
E-mail address: `dongseok@knu.ac.kr`